

Ibuprofen Dosing*

For infants and children OLDER than 6 months of age.
Give every 6-8 hours as needed for fever or pain.



At Presbyterian/St. Luke's Medical Center
The Medical Center of Aurora
North Suburban Medical Center
Rose Medical Center
Sky Ridge Medical Center
Swedish Medical Center



Weight (pounds)	Weight (kilogram)	Children's Liquid (100mg/5 mL)	Junior tablets (100 mg)	Adult tablets (200 mg)
12-21 lbs	5.5-9.9 kg	2.5 mL (½ teaspoon)	-----	-----
22-33 lbs	10-14.9 kg	5 mL (1 teaspoon)	1 tablet (100mg)	-----
34-43 lbs	15-19.9 kg	7.5 mL (1 ½ teaspoons)	1 tablet (100mg)	-----
44-55 lbs	20-24.9 kg	10 mL (2 teaspoons)	2 tablets (200mg)	1 tablet (200mg)
56-66 lbs	25-29.9 kg	12.5 mL (2 ½ teaspoons)	2 tablets (200mg)	1 tablet (200mg)
67-88 lbs	30-39.9 kg	15 mL (3 teaspoons)	3 tablets (300mg)	-----
89+ lbs	40+ kg	-----	4 tablets (400mg)	2 tablets (400mg)


* For Example, Motrin® and Advil®

See other health tips at www.RockyMountainHospitalForChildren.com



Acetaminophen Dosing*

NOT recommended for babies less than 2 months old.
Give every 4-6 hours as needed for fever or pain.



Weight (pounds)	Weight (kilogram)	Children's Liquid (160mg/5mL)	Children's Dissolve Tabs (80mg tabs)
12-17 lbs	5.5-7.9 kg	2.5 mL (½ teaspoon)	----
18-23 lbs	8-10.9 kg	3.7 mL (¾ teaspoon)	----
24-35 lbs	11-15.9 kg	5 mL (1 teaspoon)	2 tabs (160 mg)
36-47 lbs	16-20.9 kg	7.5 mL (1 ½ teaspoons)	3 tabs (240 mg)
48-59 lbs	21-26.5 kg	10 mL (2 teaspoons)	4 tabs (320 mg)
60-71 lbs	26.6-32.2 kg	12.5 mL (2 ½ teaspoons)	5 tabs (400 mg)

* For Example, Tylenol®



See other health tips at www.RockyMountainHospitalForChildren.com